



Cushion Curve
1.2 pcf (19.2 g/l)
12 in (30.5 cm) Drop Height
1" (2.5 cm) to 6" (15.2 cm) Thick


1st Impact






2nd Impact



Average of 3-5 Impacts




Cushion Curve
1.2 pcf (19.2 g/l)
18 in (45.7 cm) Drop Height
1" (2.5 cm) to 6" (15.2 cm) Thick




Cushion Curve
1.2 pcf (19.2 g/l)
24 in (61.0 cm) Drop Height
1" (2.5 cm) to 6" (15.2 cm) Thick


1st Impact






2nd Impact






Average of 3-5 Impacts



Cushion Curve
1.2 pcf (19.2 g/l)
30 in (76.2 cm) Drop Height
1" (2.5 cm) to 6" (15.2 cm) Thick




Cushion Curve
1.2 pcf (19.2 g/l)
36 in (91.4 cm) Drop Height
1" (2.5 cm) to 6" (15.2 cm) Thick




Cushion Curve
1.2 pcf (19.2 g/l)
42 in (106.7 cm) Drop Height
1" (2.5 cm) to 6" (15.2 cm) Thick


1st Impact



2nd Impact



Average of 3-5 Impacts



Cushion Curve
1.2 pcf (19.2 g/l)
48 in (121.9 cm) Drop Height
1" (2.5 cm) to 6" (15.2 cm) Thick

